

# Characterisation of Restrained Domination number and Chromatic number of a Fuzzy Graph

**J.S. Sathya**

*Research Scholar, Department of Mathematics, Mother Teresa Women's University,  
Kodaikanal, Tamilnadu India*

---

## ABSTRACT

*Let  $G(V, \sigma, \mu)$  be a simple undirected fuzzy graph. A subset  $S$  of  $V$  is called a dominating set in  $G$  if every vertex in  $V-S$  is adjacent to at least one vertex in  $S$ . A subset  $S$  of  $V$  is said to be a restrained dominating set if every vertex in  $V-S$  is adjacent to atleast one vertex in  $S$  as well as adjacent to atleast one vertex in  $V-S$ . The restrained domination number of a fuzzy graph  $G(V, \sigma, \mu)$  is denoted by  $\gamma_{fr}(G)$  which is the smallest cardinality of a restrained dominating set of  $G$ . The minimum number of colours required to colour all the vertices such that adjacent vertices do not receive the same colour is the chromatic number  $\chi(G)$ . For any fuzzy graph  $G$  a complete fuzzy sub graph of  $G$  is called a clique of  $G$ . In this paper we find an upper bound for the sum of the Restrained domination and chromatic number in fuzzy graphs and characterize the corresponding extremal fuzzy graphs.*

**Keywords:** *Fuzzy Restrained Domination Number, Chromatic Number, fuzzy graph, Clique*

## 1. INTRODUCTION

Let  $G(V, \sigma, \mu)$  be simple undirected strong fuzzy graph. The degree of any vertex  $u$  in  $G$  is the number of edges incident with  $u$  and is denoted by  $d(u)$ . The minimum and maximum degree of a vertex is denoted by  $\delta(G)$  and  $\Delta(G)$  respectively,  $P_n$  denotes the path on  $n$  vertices. The vertex connectivity  $\kappa(G)$  of a fuzzy graph  $G$  is the minimum number of vertices whose removal results in a disconnected fuzzy graph. The chromatic number  $\chi$  is defined to be the minimum number of colours required to colour all the vertices such that adjacent vertices do not receive the same colour. For any fuzzy graph  $G$  a complete sub fuzzy graph of  $G$  is called a clique of  $G$ . The number of vertices in a largest clique of  $G$  is called the clique number of  $G$ .

Let  $G(V, E)$  be a simple undirected strong fuzzy graph. A subset  $S$  of  $V$  is called a dominating set in  $G$  if every vertex in  $V-S$  is adjacent to at least one vertex in  $S$ . A subset  $S$  of  $V$  is said to be a restrained dominating set if every vertex in  $V-S$  is adjacent to atleast one vertex in  $S$  as well as adjacent to atleast one vertex in  $V-S$ . The restrained domination number, denoted by  $\gamma_{fr}(G)$  is the smallest cardinality of a restrained dominating set of a fuzzy graph  $G$ . The minimum number of

colours required to colour all the vertices such that adjacent vertices do not receive the same colour is the chromatic number  $\chi(G)$ . For any fuzzy graph  $G$  a complete sub fuzzy graph of  $G$  is called a clique of  $G$ .

If  $X$  is collection of objects denoted generically by  $x$ , then a Fuzzy set  $\tilde{A}$  is  $X$  is a set of ordered pairs:  $\tilde{A} = \{(x, \mu_{\tilde{A}}(x))/x \in X\}$ ,  $\mu_{\tilde{A}}(x)$  is called the membership function of  $x$  in  $\tilde{A}$  that maps  $X$  to the membership space  $M$  (when  $M$  contains only the two points 0 and 1). Let  $E$  be the (crisp) set of nodes. A fuzzy graph is then defined by,  $\tilde{G}(x_i, x_j) = \{(x_i, x_j), \mu_{\tilde{G}}(x_i, x_j)/(x_i, x_j) \in E \times E\}$ .  $\tilde{H}(x_i, x_j)$  is a Fuzzy Sub graph of  $\tilde{G}(x_i, x_j)$  if  $\mu_{\tilde{H}}(x_i, x_j) \leq \mu_{\tilde{G}}(x_i, x_j) \forall (x_i, x_j) \in E \times E$ ,  $\tilde{H}(x_i, x_j)$  is a spanning fuzzy sub graph of  $\tilde{G}(x_i, x_j)$  if the node set of  $\tilde{H}(x_i, x_j)$  and  $\tilde{G}(x_i, x_j)$  are equal, that is if they differ only in their arc weights.

Several authors have studied the problem of obtaining an upper bound for the sum of a domination parameter and a fuzzy graph theoretic parameter and characterized the corresponding extremal fuzzy graphs. In [5], Paulraj Joseph J and Arumugam S proved that  $\gamma + k \leq p$ . In [7], Paulraj Joseph J and Arumugam S proved that  $\gamma_c(G) + \chi \leq p + 1$ . They also characterized the class of fuzzy graphs for which the upper bound is attained. They also proved similar results for  $\gamma$  and  $\gamma_t$ . In [4], Mahadevan G introduced the concept the complementary perfect domination number  $\gamma_{cp}$  and proved that  $\gamma_{cp}(G) + \chi \leq 2n - 2$ , and characterized the corresponding external fuzzy graphs. In [9], S.Vimala and J.S.Sathya proved that  $\gamma_t(G) + \chi(G) = 2n - 5$ . They also characterised the class of fuzzy graphs for which the upper bound is attained. In this paper we obtain sharp upper bound for the sum of the restrained domination number and chromatic number and characterize the corresponding extremal fuzzy graphs. We use the following previous results.

**Theorem 1.1 [1]:** For any connected fuzzy graph  $G$ ,  $\gamma_{fr}(G) \leq n$

**Theorem 1.2 [2]:** For any connected fuzzy graph  $G$ ,  $\chi(G) \leq \Delta(G) + 1$ .

## 2. MAIN RESULTS

**Theorem 2.1:** For any connected fuzzy graph  $G$ ,  $\gamma_{fr}(G) + \chi(G) \leq 2n$  and the equality holds if and only if  $G \cong K_1$

**Proof:**  $\gamma_{fr}(G) + \chi(G) \leq n + \Delta + 1 = n + (n - 1) + 1 \leq 2n$ . If  $\gamma_{fr}(G) + \chi(G) = 2n$  the only possible case is  $\gamma_{fr}(G) = n$  and  $\chi(G) = n$ , Since  $\chi(G) = n$ ,  $G = K_n$ , But for  $K_n$ ,  $\gamma_{fr}(G) = 1$ , so that  $G \cong K_1$ . Conversely if  $G$  is isomorphic to  $K_1$ , then for  $K_1$ ,  $\gamma_{fr}(G) = 1$ , and  $\chi(G) = 1$ ,  $\gamma_{fr}(G) + \chi(G) = 2$ . Hence the proof.

**Theorem 2.2:** For any connected fuzzy graph  $G$ ,  $\gamma_{fr}(G) + \chi(G) = 2n - 1$  and the equality holds if and only if  $G \cong K_2$

**Proof:** If  $G$  is isomorphic to  $K_2$ , then for  $K_2$ ,  $\gamma_{fr}(G) = 1$ , and  $\chi(G) = 2$ .  $\gamma_{fr}(G) + \chi(G) = 2n - 1 = 3$ . Conversely assume that  $\gamma_{fr}(G) + \chi(G) = 2n - 1$ . This is possible only if  $\gamma_{fr}(G) = n$  and  $\chi(G) = n - 1$  (or)  $\gamma_{fr}(G) = n - 1$  and  $\chi(G) = n$ .

**Case (i)** Let  $\gamma_{fr}(G) = n$  and  $\chi(G) = n - 1$ .

Since  $\chi(G) = n - 1$ ,  $G$  contains a clique  $K$  on  $n - 1$  vertices. Let  $x$  be a vertex of  $G - K_{n-1}$ . Since  $G$  is connected the vertex  $x$  is adjacent to one vertex  $u_i$  for some  $i$  in  $K_{n-1}$ .  $\{x, u_i\}$  is  $\gamma_{fr}$ -set, so that  $\gamma_{fr}(G) = 2$ , we have  $n = 2$ . Then  $\chi = 1$ , which is for totally disconnected fuzzy graph. Which is a contradiction. Hence no fuzzy graph exists.

**Case (ii)** Let  $\gamma_{fr}(G) = n - 1$  and  $\chi(G) = n$

Since  $\chi(G) = n$ ,  $G = K_n$ , But for  $K_n$ ,  $\gamma_{fr}(G) = 1$ , so that  $n = 2$ ,  $\chi = 2$  Hence  $G \cong K_2$ .

**Theorem 2.3:** For any connected fuzzy graph  $G$ ,  $\gamma_{fr}(G) + \chi(G) = 2n - 2$  and the equality holds if and only if  $G \cong K_3, P_3$

**Proof:** Let  $G$  be isomorphic to  $K_3$ , then for  $K_3$ ,  $\gamma_{fr}(G) = 1$ , and  $\chi(G) = 3$ .  $\gamma_{fr}(G) + \chi(G) = 2n - 2 = 4$ . And if  $G$  is isomorphic to  $P_3$ , then for  $P_3$ ,  $\gamma_{fr}(G) = 2$ , and  $\chi(G) = 2$ .  $\gamma_{fr}(G) + \chi(G) = 2n - 2 = 4$ . Conversely assume that  $\gamma_{fr}(G) + \chi(G) = 2n - 2$ . This is possible only if  $\gamma_{fr}(G) = n$  and  $\chi(G) = n - 2$  (or)  $\gamma_{fr}(G) = n - 1$  and  $\chi(G) = n - 1$  (or)  $\gamma_{fr}(G) = n - 2$  and  $\chi(G) = n$ .

**Case (i)** Let  $\gamma_{fr}(G) = n$  and  $\chi(G) = n - 2$ .

Since  $\chi(G) = n - 2$ ,  $G$  contains a clique  $K$  on  $n - 2$  vertices. Let  $S = \{x, y\} \in G - K_{n-2}$ . Then  $\langle S \rangle = K_2$  or  $\overline{K_2}$

**Subcase (a)** Let  $\langle S \rangle = K_2$  Since  $G$  is connected,  $x$  is adjacent to some  $u_i$  of  $K_{n-2}$ . Then  $\{y, u_i\}$  for some  $i \neq j$  is  $\gamma_{fr}$ -set, so that  $\gamma_{fr}(G) = 2$  and hence  $n = 2$ . But  $\chi(G) = n - 2 = 0$ . Which is a contradiction. Hence no fuzzy graph exists.

**Subcase (b)** Let  $\langle S \rangle = \overline{K_2}$ . Since  $G$  is connected,  $x$  is adjacent to some  $u_i$  of  $K_{n-2}$ . Then  $y$  is adjacent to the same  $u_i$  of  $K_{n-2}$ . Then  $\{x, y, u_i\}$   $\gamma_{fr}$ -set, so that  $\gamma_{fr}(G)=3$  and hence  $n=3$ . But  $\chi(G)=n-2=1$  which is for totally disconnected fuzzy graph. Which is a contradiction. Hence no fuzzy graph exists, (or)  $y$  is adjacent to  $u_j$  of  $K_{n-2}$  for  $i \neq j$ . In this case  $\{x, y, u_i\}$   $\gamma_{fr}$ -set, so that  $\gamma_{fr}(G)=3$  and hence  $n=3$ . But  $\chi(G)=n-2=1$  which is for totally disconnected fuzzy graph. Which is a contradiction. Hence no fuzzy graph exists.

**Case (ii)** Let  $\gamma_{fr}(G)=n-1$  and  $\chi(G)=n-1$ .

Since  $\chi(G)=n-1$ ,  $G$  contains a clique  $K$  on  $n-1$  vertices. Let  $x$  be a vertex of  $G-K_{n-1}$ . Since  $G$  is connected,  $x$  is adjacent to one vertex  $u_i$  for some  $i$  in  $K_{n-1}$ , so that  $\{x, u_i\}$  is  $\gamma_{fr}$ -set, so that  $\gamma_{fr}(G)=2$ , we have  $n=3$ . Then  $\chi = 2$ , Hence  $G \cong P_3$

**Case (iii)** Let  $\gamma_{fr}(G)=n-2$  and  $\chi(G)=n$

Since  $\chi(G)=n$ ,  $G=K_n$ , But for  $K_n$ ,  $\gamma_{fr}(G) = 1$ , so that  $n=3$ ,  $\chi = 3$  Hence  $G \cong K_3$ . Hence the proof.

**Theorem 2.4:** For any connected fuzzy graph  $G$ ,  $\gamma_{fr}(G) + \chi(G) = 2n-3$  and the equality holds if and only if  $G \cong K_{1,3}, C_3(P_2), K_4$

**Proof:** Let  $G$  be isomorphic to  $K_{1,3}$ , then for  $K_{1,3}$ ,  $\gamma_{fr}(G) = 3$ , and  $\chi(G) = 2$ .  $\gamma_{fr}(G) + \chi(G) = 2n-3=5$ . Let  $G$  be isomorphic to  $C_3(P_2)$ , then for  $C_3(P_2)$ ,  $\gamma_{fr}(G) = 2$ , and  $\chi(G) = 3$ .  $\gamma_{fr}(G) + \chi(G) = 2n-3=5$ . Let  $G$  be isomorphic to  $K_4$ , then for  $K_4$ ,  $\gamma_{fr}(G) = 1$ , and  $\chi(G) = 4$ .  $\gamma_{fr}(G) + \chi(G) = 2n-3=5$ . Conversely assume that  $\gamma_{fr}(G) + \chi(G) = 2n-3$ . This is possible only if  $\gamma_{fr}(G)=n$  and  $\chi(G)=n-3$  (or)  $\gamma_{fr}(G)=n-1$  and  $\chi(G)=n-2$  (or)  $\gamma_{fr}(G)=n-2$  and  $\chi(G)=n-1$  (or)  $\gamma_{fr}(G)=n-3$  and  $\chi(G)=n$ .

**Case (i)** Let  $\gamma_{fr}(G)=n$  and  $\chi(G)=n-3$ .

Since  $\chi(G)=n-3$ ,  $G$  contains a clique  $K$  on  $n-3$  vertices. Let  $S=\{x, y, z\} \in G-K_{n-3}$ . Then  $\langle S \rangle = K_3, \overline{K_3}, K_2 \cup K_1, P_3$

**Subcase (i)** Let  $\langle S \rangle = K_3$ . Since  $G$  is connected,  $x$  is adjacent to some  $u_i$  of  $K_{n-3}$ . Then  $\{x, u_i\}$  is  $\gamma_{fr}$ -set, so that  $\gamma_{fr}(G)=2$  and hence  $n=2$ . But  $\chi(G)=n-3=\text{negative value}$ . Which is a contradiction. Hence no fuzzy graph exists.

**Subcase (ii)** Let  $\langle S \rangle = \overline{K_3}$  Since  $G$  is connected, one of the vertices of  $K_{n-3}$  say  $u_i$  is adjacent to all the vertices of  $S$  or two vertices of  $S$  or one vertex of  $S$ . If  $u_i$  for some  $i$  is adjacent to all the vertices of  $S$ , then  $\{x, y, z, u_i\}$  is a  $\gamma_{fr}$ -set of  $G$ , so that  $\gamma_{fr}(G)=4$  and hence  $n=4$ . But  $\chi(G)=4-3=1$  which is for totally disconnected fuzzy graph. Which is a contradiction. Hence no fuzzy graph exists. Since  $G$  is connected  $u_i$  for some  $i$  is adjacent to two vertices of  $S$  say  $x$  and  $y$  and  $z$  is adjacent to  $u_j$  for  $i \neq j$  in  $K_{n-3}$ , then  $\{x, y, z, u_j\}$  for  $i \neq j$  in  $K_{n-3}$  is  $\gamma_{fr}$ -set of  $G$ , so that  $\gamma_{fr}(G)=4$  and hence  $n=4$ . But  $\chi(G)=n-3=1$  which is totally disconnected fuzzy graph. Which is a contradiction. Hence no fuzzy graph exists. If  $u_i$  for some  $i$  is adjacent to  $x$  and  $u_j$  is adjacent to  $y$  and  $u_k$  is adjacent to  $z$ , for  $i \neq j \neq k$  in  $K_{n-3}$  then  $\{x, y, z, u_k\}$  is a  $\gamma_{fr}$ -set of  $G$ . so that  $\gamma_{fr}(G)=4$  and hence  $n=4$ . But  $\chi(G)=n-3=1$  which is for totally disconnected fuzzy graph. Which is a contradiction. Hence no fuzzy graph exists.

**Subcase (iii)** Let  $\langle S \rangle = P_3 = \{x, y, z\}$ . Since  $G$  is connected,  $x$  (or equivalently  $z$ ) is adjacent to  $u_i$  for some  $i$  in  $K_{n-3}$ . Then  $\{z, u_i\}$  is a  $\gamma_{fr}$ -set of  $G$ . so that  $\gamma_{fr}(G)=2$  and hence  $n=2$ . But  $\chi(G)=n-3=\text{negative value}$ . Which is a contradiction. Hence no fuzzy graph exists. If  $u_i$  is adjacent to  $y$  then  $\{x, z, u_j\}$  for some  $i \neq j$  is a  $\gamma_{fr}$ -set of  $G$ . so that  $\gamma_{fr}(G)=3$  and hence  $n=3$ . But  $\chi(G)=n-3=0$ . Which is a contradiction. Hence no fuzzy graph exists.

**Subcase (iv)** Let  $\langle S \rangle = K_2 \cup K_1$  Let  $xy$  be the edge and  $z$  be the isolated vertex of  $K_2 \cup K_1$  Since  $G$  is connected, there exists a  $u_i$  in  $K_{n-3}$  is adjacent to  $x$  and  $z$ . Then  $\{y, z, u_j\}$  for some  $i \neq j$  is a  $\gamma_{fr}$ -set of  $G$ , so that  $\gamma_{fr}(G)=3$  and hence  $n=3$ . But  $\chi(G)=n-3=0$ . Which is a contradiction. Hence no fuzzy graph exists. If  $z$  is adjacent to  $u_j$  for some  $i \neq j$  then  $\{y, z, u_j\}$  for some  $i \neq j$  is a  $\gamma_{fr}$ -set of  $G$ , so that  $\gamma_{fr}(G)=3$  and hence  $n=3$ . But  $\chi(G)=n-3=0$ . Which is a contradiction. Hence no fuzzy graph exists.

**Case (ii)** Let  $\gamma_{fr}(G)=n-1$  and  $\chi(G)=n-2$ .

Since  $\chi(G)=n-2$ ,  $G$  contains a clique  $K$  on  $n-2$  vertices. Let  $S=\{x, y\} \in G-K_{n-2}$ . Then  $\langle S \rangle = K_2$  or  $\overline{K_2}$

**Subcase (a)** Let  $\langle S \rangle = K_2$  Since  $G$  is connected,  $x$  (or equivalently  $y$ ) is adjacent to some  $u_i$  of  $K_{n-2}$ . Then  $\{y, u_j\}$  for some  $i \neq j$  is  $\gamma_{fr}$ -set, so that  $\gamma_{fr}(G)=2$  and hence  $n=3$ . But  $\chi(G)=n-2=1$  for which  $G$  is totally disconnected, which is a contradiction. Hence no fuzzy graph exists.

**Subcase (b)** Let  $\langle S \rangle = \overline{K_2}$  Since  $G$  is connected,  $x$  is adjacent to some  $u_i$  of  $K_{n-2}$ . Then  $y$  is adjacent to the same  $u_i$  of  $K_{n-2}$ . Then  $\{x, y, u_j\}$  for some  $i \neq j$  is  $\gamma_{fr}$ -set, so that  $\gamma_{fr}(G)=3$  and hence

$n=4$ . But  $\chi(G)=n-2=2$ . Then  $G$  is isomorphic to  $K_{1,3}$ . Otherwise  $x$  is adjacent to  $u_i$  of  $K_{n-2}$  for some  $i$  and  $y$  is adjacent to  $u_j$  of  $K_{n-2}$  for  $i \neq j$ . Then  $\{x, y, u_k\}$  for some  $i \neq j \neq k$  is  $\gamma_{fr}$ -set, so that  $\gamma_{fr}(G)=3$  and hence  $n=4$ . But  $\chi(G)=n-2=2$ . Then  $K_{n-2}=K_2$  in  $K_2$  the vertex  $u_k$  cannot be exist. Which is a contradiction. In this case also no fuzzy graph exists.

**Case (iii)** Let  $\gamma_{fr}(G)=n-2$  and  $\chi(G)=n-1$ .

Since  $\chi(G)=n-1$ ,  $G$  contains a clique  $K$  on  $n-1$  vertices. Let  $x$  be a vertex of  $K_{n-1}$ . Since  $G$  is connected the vertex  $x$  is adjacent to one vertex  $u_i$  for some  $i$  in  $K_{n-1}$  so that  $\{x, u_i\}$   $\gamma_{fr}$ -set of  $G$   $\gamma_{fr}(G)=2$ , we have  $n=4$  and  $\chi = 3$ . Then  $K=K_3$ . If  $x$  is adjacent to  $u_i$ , then  $G \cong C_3(P_2)$ .

**Case (iv)** Let  $\gamma_{fr}(G)=n-3$  and  $\chi(G)=n$

Since  $\chi(G)=n$ ,  $G=K_n$ . But for  $K_n$ ,  $\gamma_{fr}(G)=1$ , so that  $n=4$ ,  $\chi = 4$  Hence  $G \cong K_4$ . Hence the proof.

**Theorem 2.5:** For any connected fuzzy graph  $G$ ,  $\gamma_{fr}(G)+\chi(G)=2n-4$  and the equality holds if and only if  $G \cong K_{1,4}$ ,  $S(K_{1,3})$ ,  $K_3(2P_2)$ ,  $K_3(P_2, P_2, 0)$ ,  $K_4(P_1) P_4, K_5$ .

**Proof:** If  $G$  is any one of the fuzzy graphs in the theorem, then it can be verified that  $\gamma_{fr}(G)+\chi(G)=2n-4$ . Conversely assume that  $\gamma_{fr}(G)+\chi(G)=2n-4$ . This is possible only if  $\gamma_{fr}(G)=n$  and  $\chi(G)=n-4$  (or)  $\gamma_{fr}(G)=n-1$  and  $\chi(G)=n-3$  (or)  $\gamma_{fr}(G)=n-2$  and  $\chi(G)=n-2$  (or)  $\gamma_{fr}(G)=n-3$  and  $\chi(G)=n-1$  (or)  $\gamma_{fr}(G)=n-4$  and  $\chi(G)=n$ .

**Case (i)** Let  $\gamma_{fr}(G)=n$  and  $\chi(G)=n-4$ .

Since  $\chi(G)=n-4$ ,  $G$  contains a clique  $K$  on  $n-4$  vertices. Let  $S = \{v_1, v_2, v_3, v_4\} \in G - K_{n-4}$ . Then the induced subfuzzy graph  $\langle S \rangle$  has the following possible cases  $K_4, \bar{K}_4, P_4, C_4, P_3UK_1, K_2UK_2, K_3UK_1, K_{1,3}, K_4-e, C_3(1,0,0), K_2U\bar{K}_2$

In all the above cases, it can be verified that no new fuzzy graphs exists.

**Case(ii)** Let  $\gamma_{fr}(G)=n-1$  and  $\chi(G)=n-3$ .

Since  $\chi(G)=n-3$ ,  $G$  contains a clique  $K$  on  $n-3$  vertices. Let  $S=\{x, y, z\} \in G - K_{n-3}$ . Then  $\langle S \rangle = K_3, \bar{K}_3, K_2UK_1, P_3$

**Subcase (i)** Let  $\langle S \rangle = K_3$ . Since  $G$  is connected,  $x$  is adjacent to some  $u_i$  of  $K_{n-3}$ . Then  $\{z, u_i\}$  is  $\gamma_{fr}$ -set, so that  $\gamma_{fr}(G)=2$  and hence  $n=3$ . But  $\chi(G)=n-3=0$ . Which is a contradiction. Hence no fuzzy graph exists.

**Subcase (ii)** Let  $\langle S \rangle = \overline{K_3}$ . Since  $G$  is connected, one of the vertices of  $K_{n-3}$  say  $u_i$  is adjacent to all the vertices of  $S$  or two vertices of  $S$  or one vertex of  $S$ . If  $u_i$  for some  $i$  is adjacent to all the vertices of  $S$ , then  $\{x, y, z, u_i\}$  for some  $i \neq j$  in  $K_{n-3}$  is  $\gamma_{fr}$ -set of  $G$ . so that  $\gamma_{fr}(G)=4$  and hence  $n=5$ . But  $\chi(G)=n-3=2$ . Then  $K_{n-3}=K_2$  so  $G$  is isomorphic to  $K_{1,4}$ . If  $u_i$  for some  $i$  is adjacent to two vertices of  $S$  say  $x$  and  $y$  then  $G$  is connected,  $z$  is adjacent to  $u_j$  for  $i \neq j$  in  $K_{n-3}$ , then then  $\{x, y, z, u_j\}$  for some  $i \neq j$  in  $K_{n-3}$  is  $\gamma_{fr}$ -set of  $G$ . so that  $\gamma_{fr}(G)=4$  and hence  $n=5$ . But  $\chi(G)=n-3=2$ . Then  $K_{n-3}=K_2$  so  $G$  is isomorphic to  $S(K_{1,3})$ . If  $u_i$  for some  $i$  is adjacent to  $x$  and  $u_j$  is adjacent to  $y$  and  $u_k$  is adjacent to  $z$ , then  $\{x, y, z, u_l\}$  for  $i \neq j \neq k \neq l$  in  $K_{n-3}$  is  $\gamma_{fr}$ -set of  $G$ . so that  $\gamma_{fr}(G)=4$  and hence  $n=5$ .  $\chi(G)=2$  Then  $K_{n-2}=K_2$  in  $K_2$  the vertex  $u_l$  cannot be exist. Which is a contradiction. In this case also no fuzzy graph exists.

**Subcase (iii)** Let  $\langle S \rangle = P_3 = \{x, y, z\}$ . Since  $G$  is connected,  $x$ (or equivalently  $z$ ) is adjacent to  $u_i$  for some  $i$  in  $K_{n-3}$ . Then  $\{z, u_i\}$  is  $\gamma_{fr}$ -set of  $G$ . so that  $\gamma_{fr}(G)=2$  and hence  $n=3$ . But  $\chi(G)=n-3=0$ . Which is a contradiction. Hence no fuzzy graph exists. If  $u_i$  is adjacent to  $y$  then  $\{x, z, u_j\}$  is  $\gamma_{fr}$ -set of  $G$ . so that  $\gamma_{fr}(G)=3$  and hence  $n=4$ . But  $\chi(G)=n-3=1$  which is for totally disconnected fuzzy graph. Which is a contradiction. Hence no fuzzy graph exists.

**Subcase (iv)** Let  $\langle S \rangle = K_2 \cup K_1$  Let  $xy$  be the edge and  $z$  be a isolated vertex of  $K_2 \cup K_1$  Since  $G$  is connected, there exists a  $u_i$  in  $K_{n-3}$  is adjacent to  $x$  and  $z$  also adjacent to same  $u_i$  Then  $\{y, z, u_k\}$  is a  $\gamma_{fr}$ -set of  $G$ . So that  $\gamma_{fr}(G)=3$  and hence  $n=4$ . But  $\chi(G)=n-3=1$  which is for totally disconnected fuzzy graph, Which is a contradiction. Hence no fuzzy graph exists. If  $z$  is adjacent to  $u_j$  for some  $i \neq j$  then  $\{y, z, u_k\}$  is a  $\gamma_{fr}$ -set of  $G$ . So that  $\gamma_{fr}(G)=3$  and hence  $n=4$ . But  $\chi(G)=n-3=1$  which is for totally disconnected fuzzy graph, Which is a contradiction. Hence no fuzzy graph exists.

**Case (iii)** Let  $\gamma_{fr}(G)=n-2$  and  $\chi(G)=n-2$ .

Since  $\chi(G)=n-2$ ,  $G$  contains a clique  $K$  on  $n-2$  vertices. Let  $S=\{x, y\} \in G-K_{n-2}$ . Then  $\langle S \rangle = K_2$  or  $\overline{K_2}$

**Subcase (a)** Let  $\langle S \rangle = K_2$ . Since  $G$  is connected,  $x$ (or equivalently  $y$ ) is adjacent to some  $u_i$  of  $K_{n-2}$ . Then  $\{y, u_j\}$  is  $\gamma_{fr}$ -set, so that  $\gamma_{fr}(G)=2$  and hence  $n=4$ . But  $\chi(G)=n-2=2$ . Then  $G \cong P_4$ .

**Subcase (b)** Let  $\langle S \rangle = \overline{K_2}$ , since  $G$  is connected,  $x$  is adjacent to some  $u_i$  of  $K_{n-2}$ . Then  $y$  is adjacent to the same  $u_i$  of  $K_{n-2}$ . Then  $\{x, y, u_i\}$  is  $\gamma_{fr}$ -set, so that  $\gamma_{fr}(G)=3$  and hence  $n=5$ . But  $\chi(G)=n-2=3$ . So that  $K_{n-2}=K_3$  Then  $G \cong K_3(2P_2)$ , or  $y$  is adjacent to  $u_j$  of  $K_{n-2}$  for  $i \neq j$ . In this  $\{x, u_j, u_k\}$  is  $\gamma_{fr}$ -set, so that  $\gamma_{fr}(G)=3$  and hence  $n=5$ . But  $\chi(G)=3$ . So that  $K_{n-2}=K_3$  Then  $G \cong K_3(P_2, P_2, 0)$

**Case (iv)** Let  $\gamma_{fr}(G)=n-3$  and  $\chi(G)=n-1$ .

Since  $\chi(G)=n-1$ ,  $G$  contains a clique  $K$  on  $n-1$  vertices. Let  $x$  be a vertex of  $G-K_{n-1}$ . Since  $G$  is connected the vertex  $x$  is adjacent to one vertex  $u_i$  for some  $i$  in  $K_{n-1}$ , then  $\{x, u_i\}$  is  $\gamma_{fr}$ -set of  $G$  so that  $\gamma_{fr}(G)=2$ , we have  $n=5$  and  $\chi=4$ . Then  $K_{n-1}=K_4$  Let  $u_1, u_2, u_3, u_4$  be the vertices of  $K_4$ . Then  $x$  must be adjacent to only one vertex of  $G-K_3$ . Without loss of generality let  $x$  be adjacent to  $u_1$ . If  $d(x)=1$ , then  $G \cong K_4(P_2)$ .

**Case (v)** Let  $\gamma_{fr}(G)=n-4$  and  $\chi(G)=n$

Since  $\chi(G)=n$ ,  $G=K_n$ , But for  $K_n$ ,  $\gamma_{fr}(G)=1$ , so that  $n=5$ ,  $\chi=5$ . Hence  $G \cong K_5$ . Hence the proof.

### 3. CONCLUSION

In this paper, upper bound of the sum of fuzzy restrained domination and chromatic number is proved. In future this result can be extended to various domination parameters. The structure of the graphs had been given in this paper can be used in models and networks. The authors have obtained similar results with large cases of fuzzy graphs for which  $\gamma_{fr}(G)+\chi(G)=2n-5$ ,  $\gamma_{fr}(G)+\chi(G)=2n-6$ ,  $\gamma_{fr}(G)+\chi(G)=2n-7$ ,  $\gamma_{fr}(G)+\chi(G)=2n-8$

### REFERENCES

- [1] Teresa W. Haynes, Stephen T. Hedemiemi and Peter J. Slater (1998), fundamentals of Domination in fuzzy graphs, Marcel Dekker, Newyork.
- [2] Hanary F and Teresa W. Haynes,(2000), Double Domination in fuzzy graphs, ARC Combinatoria 55, pp. 201-213
- [3] Haynes, Teresa W.(2001): Paired domination in Fuzzy graphs, Congr. Number 150
- [4] Mahadevan G, Selvam A, (2008): On independent domination number and chromatic number of a fuzzy graph, Acta Ciencia Indica, preprint
- [5] Paulraj Joseph J. and Arumugam S.(1992): Domination and connectivity in fuzzy graphs, International Journal of Management and systems, 8 No.3: 233-236.
- [6] Mahadevan G,(2005): On domination theory and related concepts in fuzzy graphs, Ph.D. thesis, Manonmaniam Sundaranar University, Tirunelveli, India.
- [7] Paulraj Joseph J. and Arumugam S.(1997): Domination and colouring in fuzzy graphs. International Journal of Management and Systems, Vol.8 No.1, 37-44.



- [8] Paulraj Joseph J, Mahadevan G, Selvam A (2004). On Complementary Perfect domination number of a fuzzy graph, *Acta Ciencia India*, vol. XXXIM, No.2,847(2006).
- [9] Vimala S, Sathya J.S, “Graphs whose sum of Chromatic number and Total domination equals to  $2n-5$  for any  $n>4$ ”, *Proceedings of the Heber International Conference on Applications of Mathematics and statistics*, Tiruchirappalli pp 375-381
- [10] Vimala S, Sathya J.S, “Total Domination Number and Chromatic Number of a Fuzzy Graph”, *International Journal of Computer Applications* (0975 – 8887) Volume 52– No.3, August 2012
- [11] Vimala S, Sathya J.S, “Connected point set domination of fuzzy graphs”, *International Journal of Mathematics and Soft Computing*, Volume-2, No-2 (2012),75-78
- [12] Vimala S, Sathya J.S, “Some results on point set domination of fuzzy graphs”, *Cybernetics and Information Technologies*, Volume 13, No 2 2013 Print ISSN: 1311-9702; Online ISSN: 1314-4081 Bulgarian Academy Of Sciences, Sofia -2013.
- [13] Vimala S, Sathya J.S, “The Global Connected Domination in Fuzzy Graphs”, *Paripex-Indian Journal of research*, Volume 12, Issue: 12, 2013, ISSN - 2250-1991
- [14] Vimala S, Sathya J.S, “Efficient Domination number and Chromatic number of a Fuzzy Graph”, *International Journal of Innovative Research in Science, Engineering and Technology*, Vol. 3, Issue 3, March 2014, ISSN: 2319-8753